



Student Allergy/Anaphylaxis Action Plan

Student
Photo

Student Name _____ D.O.B. _____ Teacher _____
 School Nurse _____ Phone Number _____
 Health Care Provider _____ Preferred Hospital _____
 History of Asthma No Yes-Higher risk for severe reaction

ALLERGY: (check appropriate) **To be completed by Health Care Provider**

- Foods (list):**
- Medications (list):**
- Latex: Circle: Type I (anaphylaxis) Type IV (contact dermatitis)**
- Stinging Insects (list):**

RECOGNITION AND TREATMENT

Chart to be completed by Health Care Provider <i>ONLY</i>		Give CHECKED Medication	
<i>If food ingested or contact w/allergen occurs:</i>		Epinephrine	Antihistamine
No symptoms noted	<input type="checkbox"/> Observe for other symptoms		
Mouth	Itching, tingling, or swelling of lips, tongue, mouth		
Skin	Hives, itchy rash, swelling of the face or extremities		
Gut+	Nausea, abdominal cramps, vomiting, diarrhea		
Throat+	Tightening of throat, hoarseness, hacking cough		
Lung+	Shortness of breath, repetitive coughing, wheezing		
Heart+	Thready pulse, low BP, fainting, pale, blueness		
Neuro+	Disorientation, dizziness, loss of consciousness		
If reaction is progressing (several of the above areas affected), GIVE:			
<i>The severity of symptoms can quickly change. + Potentially life-threatening</i>			

DOSAGE:

Epinephrine: Inject into outer thigh **0.3 mg** OR **0.15 mg**

Antihistamine: Liquid Diphenhydramine (Benadryl®) _____ mg. To be given by mouth *only if able to swallow*.

Other:

- This child has received instruction in the proper use of the Auto-injector: EpiPen® or Twinject® (circle one). It is my professional opinion that this student **SHOULD** be allowed to carry and use the auto-injector independently. The child knows when to request antihistamine and has been advised to inform a responsible adult if the auto-injector is self-administered.
- It is my professional opinion that this student **SHOULD NOT** carry the auto-injector.

Health Care Provider Signature _____ **Phone:** _____ **Date** _____

EMERGENCY CALLS

1. **Call 911.** State that an allergic reaction has been treated, and additional epinephrine may be needed.
2. Call parents/guardian to notify of reaction, treatment and student's health status.
3. Treat for shock. Prepare to do CPR.
4. Accompany student to ER if no parent/guardians are available.

Side 2: To Be Completed by Parent/Guardian, Student and School

Allergy/Anaphylaxis Action Plan *(continued)* Student Name _____ D.O.B. _____

Each school will have 2 auto-injectors and liquid Diphenhydramine (Benadryl®) available during regular school hours. If your child participates in before or after school activities, your child will need to have an auto-injector on their person.

Parent/Guardian AUTHORIZATIONS

- I want this allergy plan implemented for my child; **I want my child to carry an auto-injector** and I agree to release the school district and school personnel from all claims of liability if my child suffers any adverse reactions from self-administration of EpiPen.
- I want this plan implemented for my child and I **do not** want my child to self-administer epinephrine.

Your signature gives permission for the nurse to contact and receive additional information from your health care provider regarding the allergic condition(s) and the prescribed medication.

Parent/Guardian Signature: _____ **Phone:** _____ **Date:** _____

Student Agreement:

- I have been trained in the use of my auto-injector and allergy medication and understand the signs and symptoms for which they are given;
- I agree to carry my auto-injector with me at all times;
- I will notify a responsible adult (teacher, nurse, coach, noon duty, etc.) **IMMEDIATELY** when my auto-injector (epinephrine) is used;
- I will not share my medication with other students or leave my auto-injector unattended;
- I will not use my allergy medications for any other use than what it is prescribed for.

Student Signature: _____ **Date** _____

Back-up medication is stored at school Yes No

Approved by Nurse/Principal Signature: _____ **Date** _____

PREVENTION: Avoidance of allergen is crucial to prevent anaphylaxis.

Critical components to prevent life threatening reactions: √ Indicates activity completed by school staff

<input type="checkbox"/>	Encourage the use of Medic-Alert bracelets
<input type="checkbox"/>	Notify nurse, teacher(s), front office and kitchen staff of known allergies
<input type="checkbox"/>	Use non-latex gloves and eliminate powdered latex gloves in schools
<input type="checkbox"/>	Ask parents to provide non-latex personal supplies for latex allergic students
<input type="checkbox"/>	Post "Latex Reduced Environment" sign at entrance(s) of building
<input type="checkbox"/>	Encourage a no-peanut zone in the school cafeteria
<input type="checkbox"/>	Other:

STAFF MEMBERS TRAINED

NAME	TITLE	LOCATION/ROOM#	TRAINED BY

EMERGENCY CONTACTS

	NAME	HOME#	WORK#	CELL#
Parent/Guardian				
Parent/Guardian				
Other:				
Other:				